



UTAH FACIAL PLASTICS

Surgery • Med Spa • Hair Restoration

PRE-OPERATIVE INSTRUCTIONS NASAL SURGERY

3-4 WEEKS BEFORE SURGERY:

- **Obtain Medical Evaluation:** If instructed by UFP, visit your primary care doctor to complete your medical evaluation. This should be completed at least 2 weeks before surgery. All results can be faxed to 801-615-2258.
- **Smoking:** Tobacco, cigarettes, and nicotine consumption will cause poor wound healing, longer healing times, and excessive scarring. You must discontinue these products more than 4 weeks prior to surgery. Otherwise, the surgery will be cancelled.

2 WEEKS BEFORE SURGERY:

- **Diet/Supplement restrictions:** Avoid foods, drinks and herbal supplements that can increase the risk of bleeding and bruising. Avoid aspirin, Ibuprofen, Naproxen, Motrin, blood thinners, vitamin E, omega-3, fish oil, and alcohol. Use Tylenol in place of other over-the-counter pain medications.
- **Vitamin C:** Begin taking 1000 – 3000mg per day as tolerated 2 weeks. Vitamin C aids in wound healing.
- **Post surgery:** If your destination after surgery is over 60 minutes from the office, you must stay locally.
- **Waxing:** It is recommended to wax your nose two weeks prior to surgery.

1 WEEK BEFORE SURGERY:

- **Review Instructions with Caregiver:** Confirm who will be driving you to and from surgery.
- **Alcohol:** Do not drink alcohol 1 week before and after surgery. Alcohol consumption may raise the blood pressure and initiate bleeding.
- **False eyelashes:** Have any false lashes removed, including extensions or strip lashes.

NIGHT BEFORE SURGERY:

- **Arrival time:** Utah Facial Plastics will call the day before surgery with the arrival time. If you do not receive a call or for further questions, call the appropriate number below:
 - Utah Facial Plastics: (801)-776-2220
- **Shower:** The night or morning of surgery, take a shower and wash your hair. This will help reduce the risk of infection by eliminating bacteria.
- **Set Up Home Recovery Area:** This may include pillows, blankets, books, television, and anything else to assist with a comfortable recovery.

DAY OF SURGERY:

- **Diet:** DO NOT FAST. Eat and drink as normal. DO NOT drink any caffeine.
- **Dress Comfortably:** Dress in comfortable, clean, and loose-fitting clothes. Shirts that can be buttoned or zipped-up are preferred to avoid snagging the nose.
- **Arrival to Facility:** Have someone drive you to and from your designated surgery location.
- **Do NOT:** wear any lotion, sunscreen, mascara, makeup, gel nail polish or press-on nails, jewelry (remove all piercings), cosmetic creams.

POST-OPERATIVE INSTRUCTIONS FOR NASAL SURGERY

DAYS 1-7 AFTER SURGERY:

- **Gentle Movement is Important:** Make sure to get out of bed and slowly walk around the room with some assistance to avoid lower leg blood clots. When in bed or on the couch, make sure to move your legs and ankles periodically. Take occasional deep breaths for the first 24 hours to keep your lungs clear.
- **Cold Compress:** Use cold compress on your eyes, cheeks, and forehead (not nose), without getting your nose cast wet. Twenty minutes on and 20 minutes off. The first 3 days are the most crucial for icing.
- **Sleep:** Sleep on your back with your upper body elevated about 45-60 degrees. Try using 4-5 pillows, an airplane neck pillow & a wedge pillow) or recliner. Do not sleep on your side. **DO NOT SLEEP FLAT.** Keeping your body more upright will minimize swelling. Do this for at least 1-2 weeks or even longer. Lying flat will increase swelling and scar tissue.
- **Drip Pad:** You will have a gauze drip pad placed beneath your nose. Change this as needed for the first 72 hours (about 3 days) following rhinoplasty surgery. It is common to change this every 15 minute for the first several hours following rhinoplasty. If you completely saturate the pad with bright red blood every five minutes, please call our office.
- **Saline Spray:** Start using Saline spray the following morning. 1-2 sprays each side, every hour while awake. Continue nasal saline for 1 month following your surgery.
- **Cast:** The cast must remain on your nose for 7 days after your procedure. Keep the cast dry by carefully washing around your face with a mild washcloth or soap. Do not take a hot/steamy shower as this can cause the cast to fall off prematurely.
- **Diet:** Eat as tolerated and take small bites. Soft foods are easier to swallow since you will not be able to breathe through your nose and chewing may be uncomfortable.

NIGHT BEFORE YOUR CAST REMOVAL:

- **Shower:** Get your face and nose cast soaking wet for 20 minutes under the shower with lukewarm water (NOT HOT WATER) the night before and the day of your cast removal (2 showers) so that we can remove your cast easily and painlessly. If you have dry skin take a third shower. Let the water hit the top of your head and forehead and drip over your nose. Do not let shower spray directly hit your nose.

AFTER CAST REMOVAL (FOR 30 DAYS):

- **Exercise:** Slowly resume your activities beginning the second week. You may begin by walking or similar light exercise. Strenuous exercise may be resumed in 4-5 weeks. Build up to this level slowly. Contact and semi-contact sports need to be avoided for 3 months. Full swimming is allowed after 6 weeks.
- **Sun & Heat:** Your skin should be protected from sun exposure for at least six months after surgery. A sunburn will cause the nose to swell dramatically and delay the result. Sun avoidance or protection with a hat is preferred. You may begin wearing a zinc-based sun block once the cast is removed.
- **Flying:** Avoid unnecessary flights (vacation, etc.) for 3 months. We recommend “taping” of your nose when you fly back home if you are an out-of-town patient.

MEDICATIONS:

- **Pain:** You will be prescribed a pain medication for post-operative pain control. Good pain control reduces bleeding after surgery. If your discomfort after surgery is not strong, you can take Tylenol instead of the prescribed medication. Do not take Tylenol with the pain medication, as most often the medication you are prescribed will have Tylenol in it. Do not exceed 4,000 mg of Acetaminophen in any 24-hour period. Take medication with food to minimize the risk of nausea. Keep in mind some discomfort and pressure or throbbing is to be expected.
- **Constipation:** You may experience constipation if taking narcotic pain relievers. MiraLAX or other over-the-counter laxatives are recommended. Do not wait to take it until you are constipated. Begin taking laxatives with the first narcotic pain pill.

ACTIVITY:

- **Physical Activity:** Refrain from any physically exerting activity. No bending over, straining, or lifting more than 10 lbs. for the first week after surgery. **DO NOT RAISE YOUR BLOOD PRESSURE OR HEART RATE.** Being too active increases bleeding, swelling and scar tissue.
- **Return to Work:** Most patients return to work after their cast is removed at their one-week appointment. This, however, depends on their job responsibilities and amount of face-to-face interaction and bruising.
- **Driving:** Do NOT operate a vehicle or make important decisions until you have been off narcotic pain medications for 24 hours or more. Use good judgment.
- **Sneezing:** *If you must sneeze, do so with your mouth open to reduce pressure.*
- **Blowing Your Nose:** No nose blowing and no nose wiping for a month. This may give you a nosebleed or dislodge grafts.
- **Hot Tubs/Baths/Swimming Pools:** No tub baths or Jacuzzi for 1-2 weeks which may cause swelling, bleeding, and fainting. Do not submerge your head under public pool, lake, or ocean water for 6 months till internal incisions have healed and sealed.

WHAT TO EXPECT WITH HEALING:

- **Swelling:** Swelling is to be expected for many months following a rhinoplasty. Your nose is swollen. Your cheeks, lips, forehead and between the eyes are all swollen. It is all to be expected. Your face may look “odd” to you. This is normal. You may even experience shock, panic or crying. These are normal reactions in a minority of patients. The more you ice the cheeks, eyes, and forehead the first 2-3 days, the easier and faster your recovery will be moving forward.
- **Bruising:** It is normal to have bruising around the eyes. Most bruises disappear after about 3-4 weeks. The bruises will go from a purple/blue color to a yellow/green shade as it starts to resolve.
- **Breathing:** Internal nose swelling from the surgery can restrict airflow through the nose. One side of the nose may be easier to breathe through than the other. While some patients may get improved nasal airflow after a few weeks, some take longer.
- **Tip Numbness:** It is normal for the tip of the nose to be numb. This is temporary and will improve with time.

EMERGENCY SITUATIONS – WHEN TO CALL THE OFFICE OR GO TO THE HOSPITAL

- **Signs of Infection:** Spreading redness, worsening swelling, increased drainage or drainage of pus, worsening pain, warmth at incision site and temperature over 101°F.
- **Excessive Bleeding:** If the dressings are saturated with bright red blood and you are having to make very frequent dressing changes.
- **Other Emergency Situations:** Shortness of breath or difficulty breathing, chest pain, lightheadedness that does not quickly resolve, severe vomiting, pain, or asymmetric swelling in your legs.

Utah Facial Plastics After Hours Answering Service – (801) 299-7660

APPOINTMENT TYPE	DATE / TIME	LOCATION	STAFF/PROVIDER
Surgery: Time TBD Day before			
Post-Op Day 7: Cast Removal			
3-4 Week Follow Up:			

